

Golden redfish

Sebastes norvegicus

STOCK DESCRIPTION AND MANAGEMENT UNITS

Golden redfish (*Sebastes norvegicus*) in ICES division 5.a (Iceland), 5.b (Faroe Islands) and Subarea 14 (East Greenland) have been considered as one management unit. Catches in ICES Subarea 6 have traditionally been included in this report. Data from ICES Subarea 6 are, however, not used in the assessment.

SURVEYS

This section describes results from various surveys conducted annually on the continental shelves and slopes of ICES Subareas 5 and 14.

DIVISION 5.A

Two bottom trawl surveys are conducted in Icelandic waters, the Icelandic spring groundfish survey (spring survey) and the Icelandic autumn groundfish survey (autumn survey). The spring survey has been conducted annually in March since 1985 and the autumn survey annually in October since 1996. The autumn survey was not conducted in 2011. The calculation of the survey indices includes length dependent diel vertical migration of the species.

Two survey indices are calculated from these surveys but only the index from the spring survey is used in the assessment of golden redfish. Length disaggregated indices from the spring survey are used in the Gadget model. Age-length keys from the autumn survey in 2 cm length groups are used in the Gadget model.

The total biomass of golden redfish as observed in the spring survey decreased from 1988 to a record low in 1995 (Figure 1 and Table 1). From 2000 to 2016 the biomass increased, with some fluctuation, to the highest value in the time-series. Since then, the index has decreased and was in 2019–2022 similar as in 2014 and 2015. The CV of the survey indices has been considerably higher after 2002.

The total biomass index from the autumn survey shows a similar trend as in the spring survey, that is, has gradually increased from 2000 to the highest value in the time series in 2014. The total biomass index has since then been fluctuating around the 2014 level but decreased sharply in 2020 and 2021 (Figure 1 and Table 1).

Length disaggregated indices from the spring survey shows that the peaks in length 4–11 cm, which can be seen first in 1987 (the 1985 cohort) and then in 1991–1992 (the 1990 cohort), reached the fishable stock approximately 10 years later (Figure 2). The increase in the survey index between 1995 and 2005 reflects the recruitment of these two strong year classes. During the 1999–2008 period the abundance of small redfish was highest in 2000–2003 (Figure 1). Very little of small redfish was observed in the spring survey 2009–2020 but in recent two years the index increased (Figure 1). The recruitment index in 2022 was the highest value observed since 2000.

In recent years, the modes of the length distribution in both surveys have shifted to the right and are narrower. The abundance of golden redfish smaller than 30 cm has decreased since 2006 in both surveys and is now at the lowest level in the time-series (Figures 1-3).

Age disaggregated abundance indices from the autumn survey are shown in Figure 4 and Table 2. The sharp increase in the survey indices since 2005 reflects the recruitment of the 1996–2007 cohorts. The 1996–2002 cohorts are gradually disappearing from the stock and the 2003–2008 cohorts are now the most abundant year classes. The age disaggregated abundance indices indicate that all year classes since 2009 are small.




Figure 1: Indices of golden redfish in ICES Division 5.a (Icelandic waters) from the groundfish surveys in March 1985–2022 (blue line and shaded area) and October 1996–2021 (yellow lines and shaded areas). The shaded areas represent 95% CI.




Figure 2: Length disaggregated abundance indices (yellow area) of golden redfish from the groundfish survey in March 1985–2022 conducted in Icelandic waters. The blue line is the mean of total indices 1985–2022.




Figure 3: Length disaggregated abundance indices (yellow area) of golden redfish from the groundfish survey in October 1996–2021 conducted in Icelandic waters. The blue line is the mean of total indices 1996–2021. The survey was not conducted in 2011.




Figure 4: Age disaggregated abundance indices of golden redfish in the groundfish survey in October conducted in Icelandic waters 1996–2021. The survey was not conducted in 2011.

DIVISION 5.B

CPUE of golden redfish in Division 5.b are available from the Faroes spring groundfish survey from 1994–2022 and the summer survey 1996–2021. Both surveys show similar trends in the indices from 1998 onwards with sharp declines between 1998 and 1999 (Figure 5). Survey CPUE index in the spring survey since 2000 has been stable at low level. The CPUE index in the summer survey shows a similar trend as in the spring survey and has gradually decreased and is at the lowest level recorded in 2020 but increased in 2021.

The fish caught in the surveys in Division 5.b (Figures 6 and 7) is on the average larger than the fish caught in the Icelandic surveys (Figures 2 and 3) and the survey conducted in East Greenland waters (Figure 8). The modes of the length distribution in both surveys in Faroes waters have shifted to the right towards larger fish, and very little of fish smaller than 35 cm is caught. This is the same trend as observed in Icelandic and East Greenland waters.




Figure 5: CPUE of golden redfish in the Faroes spring groundfish survey 1994–2022 (blue line) and the summer groundfish survey 1996–2021 (red line) in ICES Division 5.b.




Figure 6 Length distribution (yellow area) of golden redfish in the Faeroes spring groundfish survey 1994–2022. The blue line is the mean for 1994–2022.




Figure 7 Length distribution (yellow area) of golden redfish in the Faeroes summer groundfish survey 1996–2021. The blue line is the mean for 1996–2021.

SUBAREA 14

The German groundfish survey has been conducted annually in the autumn from 1982 to 2017 and in 2019–2020 covering shelf areas and the continental slopes off West and East Greenland. The survey was not conducted in 2018 and 2021 because of various factor such as research vessel breakdown, bad weather and the Covid-19 pandemic.

Relative abundance and biomass indices for golden redfish (fish > 17 cm) from the German groundfish survey are illustrated in Figure 8. After a severe depletion of the golden redfish stock on the traditional fishing grounds around East Greenland in the early 1990s, the survey estimates showed a significant increase from 2003, both in biomass and abundance (Figure 8). The survey indices in 2007–2020 were high but fluctuated. The biomass survey index in 2014–2016 were at the highest level in the time-series but decreased in 2017–2020 to a similar level as in 2006 (Figure 8a). It should be noted that the CV for the indices is high, and the increase is driven by few very large hauls. In 2010–2020, the biomass of pre-fishery recruits (17–30 cm) has decreased gradually compared to previous five years and in 2017–2020 very little of 17–30 cm fish was observed (Figure 8c).

Abundance indices of redfish smaller than 18 cm from the German groundfish survey show that juveniles were abundant in 1993 and 1995–1998. Since 2008, the survey index has been very low and in recent years at the lowest value recorded since 1982. Juvenile redfish were only classified to the genus *Sebastes* spp., as species identification of small specimens is difficult due to very similar morphological features. The 1999–2020 survey results indicate low abundance and are like those observed in the late 1980s. The Greenland shrimp and fish shallow water survey (no survey conducted 2017–2019) also shows that very few juvenile redfish (< 18 cm, not classified to species) were present.




Figure 8: Golden redfish (> 17 cm). Survey indices for East Greenland (ICES Subarea 14) from the German groundfish survey 1985–2020. a) Total biomass index, b) total abundance index, c) biomass index divided by size classes (17–30 cm and >30 cm). The survey was not conducted in 2018 and 2021.




Figure 9. Golden redfish (>17 cm). Length frequencies for East Greenland (ICES Subarea 14) 1982–2020. The survey was not conducted in 2018 and 2021.

FISHERY

LANDINGS

Total landings of golden redfish gradually decreased by more than 70% in 1982–1994 or from 130 429 t in 1982 to 43 515 t in 1994 (Table 3 and Figure 10). Since then, total annual landings have varied between 33 451 and 59 698 t. The total landings in 2021 were 43 426 t, which is 2 771 t less than in 2020. Most of the golden redfish catch or 90–98% has been taken in ICES Division 5.a.

Landings of golden redfish in Division 5.a declined from 97 899 t in 1982 to 38 669 t in 1994 (Table 3). Since then, landings have varied between 31 686 t and 54 041 t, highest in 2016. The annual landings since 2016 have decreased and were 39 616 t in 2021, 1 072 t less than in 2020. The landings for the 2020/2021 fishing year were 18% higher than allocated quota of 34 379 t. The reasons for the implementation errors are related to the management system that allow for transfers of quota share between fishing years and conversion of TAC from one species to another.

Between 90–95% of the golden redfish catch in Division 5.a is taken by bottom trawlers targeting redfish (both fresh fish and factory trawlers; vessel length 48–65 m). The remaining catches are partly caught as bycatch in gillnet, long-line, and lobster fishery. In 2021, as in previous years, most of the catches were taken along the shelf southwest, west, and northwest of Iceland (Figure 11). Higher proportion of the catches is now taken along the shelf northwest of Iceland and less south and southwest.

In Division 5.b (Faroese waters), annual landings decreased from 9194 t in 1985 to less than 700 t in the 2006–2016 period (Table 3). In 2017, landings increased to 1397 t, the highest landings since 2005. The landings in 2021 decreased to 178 t, 1126 t less than in 2020 and similar as in 2016. Most of the golden redfish caught in Division 5.b is taken by pair and single trawlers (vessels larger than 1000 HP).

In Subarea 14 (East Greenland waters), the landings of golden redfish reached a record high of 30 962 t in 1982 but decreased drastically within the next three years to 2117 t in 1985 (Figure 10 and Table 3). During the period 1985–1994, the annual landings from Subarea 14 varied between 687 and 4255 t. There was little or no direct fishery for golden redfish from 1995 to 2009 and landings were 200 t or less, mainly taken as bycatch in the shrimp fishery. In 2010, landings of golden redfish increased considerably and were 1650 t, mainly due to increased *S. mentella* fishery in the area. Annual landings 2010–2015 have been between 1000 t and 2700 t but increased to 5442 t in 2016 which is the highest landings since 1983. The landings in 2021 were 3 532 t, 573 t less than in 2020.

Annual landings from Subarea 6 increased from 1978 to 1987 followed by a gradual decrease to 1992 (Table 3). From 1995 to 2004, annual landings have ranged between 400 and 800 t, but decreased to 137 t in 2005. Little or no landings of golden redfish were reported from Subarea 6 in 2006–2021 and were 100 t in 2021.




Figure 10. Nominal landings of golden redfish in tonnes by ICES Subareas and Divisions 1978–2021.




Figure 11. Geographical distribution of golden redfish bottom trawl catches in ICES Division 5.a 2006–2021.

DISCARD

Comparison of sea and port samples from the Icelandic discard sampling program does not indicate significant discarding due to high grading (Pálsson et al. 2010), possibly due to area closures of important nursery grounds west off Iceland. Substantial discard of small redfish took place in the deep-water shrimp fishery from 1986 to 1992, before sorting grids became mandatory. Since then, the discard has been insignificant both due to the sorting grid and much less abundance of small redfish in the region.

Discard of redfish species in the shrimp fishery in ICES Division 14.b is currently considered insignificant.

BIOLOGICAL DATA FROM THE COMMERCIAL FISHERY

The table below shows sampling from bottom trawl catches by ICES divisions in 2021. No sampling of the commercial catch from Subarea 6 was carried out.

Sampling from the bottom trawl catches in Icelandic waters (5.a) is considered sufficient and covers the spatial distribution of catches. The sampling coverage in 2021 is shown in Figure 12.

Area	Nation	Gear	Landings (t)	Samples	No. length measured	No. Age read
5.a	Iceland	Bottom trawl	40 688	65	9 191	834
5.b	Faroe Islands	Bottom trawl	187			
14	Greenland	Bottom trawl	3 532			




Figure 12. Demersal trawl fishing grounds in 2021 as reported in logbooks and positions of samples of golden redfish taken from landings (crosses).

LANDINGS BY LENGTH AND AGE

Length distributions from the Icelandic commercial trawler fleet in 1976–2021 show that most of the fish caught is 30–45 cm (Figure 13). The modes of the length distributions range between 35 and 40 cm and has over the past decade shifted to the right. The length distributions in 2012–2021 are narrower than previously, with less than average of small fish (<35 cm), and the mean length has increased by almost 5 cm.

Catch-at-age data from the Icelandic fishery in Division 5.a show that the 1985 cohort dominated the catches from 1995–2002 and the 1990 cohort dominated the catches in 2003–2007 (Figure 14 and Table 4). The 2004–2009 cohorts (ages 12–17) were the most dominant year classes in the fishery in 2021. There is a substantial decrease of 7–10-year-old fish in the catch, compared to recent previous years, an additional indicator of low recruitment in recent years, as observed in all surveys conducted in East Greenland and Icelandic waters.

The average total mortality (Z), estimated from the 25-year series of catch-at-age data (Figure 15) is about 0.20 for age 13 years and older.

Length distribution from the Faroese commercial catches 2001–2019 shows that the fish caught are on average larger than 40 cm with modes between 45 cm and 50 cm (Figure 16).

No length data from the catches have been available for several years from East Greenland waters (Subarea 14).




Figure 13. Length distribution (blue shaded area) of golden redfish in Icelandic waters (ICES Division 5.a) in the commercial landings of the Icelandic bottom trawl fleet 1976–2021. The yellow line is the mean of the years 1976–2021.




Figure 14. Catch-at-age of golden redfish in numbers in ICES Division 5.a 1995–2021. Bar size is indicative of the catch in numbers and bars are colored by cohort.




Figure 15. Catch curve of the 1981–2005 cohorts of golden redfish based on the catch-at-age data in ICES Division 5.a 1995–2021.




Figure 16. Length distribution of golden redfish from Faroese catches in ICES Division 5.b in 2001-2019.

CATCH PER UNIT EFFORT

The un-standardized CPUE index from the Icelandic bottom trawl fleet operating in Division 5.a has increased sharply from 2006 to the highest level in the time series in 2017-2019 (Figure 17). CPUE has since then decreased although it remains high. Effort towards golden redfish has since 1986 gradually decreased and is at the lowest level recorded (Figure 17). CPUE derived from logbooks is not considered indicative of stock trends, however the information contained in the logbooks on effort, spatial and temporal distribution of the fishery is of value.

CPUE from other areas are not available. This is because no separation of *S. norvegicus*/*S. mentella* is made in the catches.




Figure 17. CPUE of golden redfish from Icelandic trawlers 1978–2021 where golden redfish catch composed at least 50% of the total catch in each haul (black line), 80% of the total catch (red line), and in all tows where golden redfish was caught (blue line).

ANALYTICAL ASSESSMENT

The stock was benchmarked in January 2014 and a management plan evaluated and adopted (WKREDMP, ICES 2014). The benchmark group agreed to base the advice on the Gadget framework (see <http://www.hafro.is/gadget> for further details).

GADGET MODEL

DATA AND MODEL SETTINGS

Below is a brief description of the data used in the model and model settings is given.

Data used in the Gadget model are:

- Length disaggregated survey indices 19–54 cm in 2-cm length increments from the Icelandic groundfish survey in March 1985–2022 and the German survey in East Greenland 1984–2021. The German survey index in 2018 (survey not conducted) is based on the average of the 2017 and 2019 and the 2021 (survey not conducted) index is set as the same as in 2020.
- Survey indices are combined (Figure 18) and the German survey gets half the weight compared to what is presented in Figure 8. This was done to avoid extrapolation to areas not surveyed, and hence reduce noise. By using the stratification used to calculate indices, each station in the German survey would get 2.5 times more weight compared to the Icelandic survey.
- Length distributions from the Icelandic (1972–2022), Faroe Islands (1980–2020) and East Greenland (1975–2004) commercial catches.
- Landings by 6-month period from Iceland, Faroe Islands and East Greenland.
- Age-length keys and mean length at age from the Icelandic groundfish survey in October 1996–2021.
- Age-length keys and mean length at age from the Icelandic commercial catch 1995–2021.

Model settings:

- The simulation period is from 1970 to 2027 using data until the first half of 2022 for estimation. Two time-steps are used each year. The ages used were 5 to 30 years, where the oldest age is treated as a plus group (fish 30 years and older).
- Modelled length ranged between 19–54 cm.
- Commercial catches are split by country and implemented as separate fleets. Survey catch distribution data are modelled as a separate fleet.
- Recruitment was set at age 5.

Estimated parameters are:

- Number of fishes when the simulation starts (8 parameters).
- Recruitment at age 5 each year (54 parameters).
- Length at recruitment (3 parameters).
- Parameters in the growth equation; (2 parameters).
- Parameter β of the beta-binomial distribution controlling the spread of the length distribution.
- Selection pattern of the three commercial fleets assuming logistic selection (S-shape) (3x2 parameters).
- Selection pattern of the survey fleet assuming an Andersen selection curve (bell-shape) (3 parameters).

It should be noted that the length disaggregated indices are from the spring survey, but the age data are from the autumn survey conducted six months later. The surveys could have different catchability,

but the age data are used as proportions within each 2 cm length group, so it should not have an impact on the results. Growth in between March and October is included in the model.

Assumptions done in the predictions:

- Recruitment at age 5 in 2023 and onwards was set as the average of the five smallest estimated year classes 1980–2007 or 39.5 million. The reason is indication of poor recruitment in recent years, but estimated recruitment was even lower.
- Catches in 2022 were set as the sum of expected landings, accounting for interannual transfer from 2021. Catches in 2022 were set as the sum of expected landings, accounting for interannual transfer from 2021.
- The estimated selection pattern from the Icelandic fleet was used for projections.




Figure 18. Biomass index from Iceland (blue) and Greenland (red), based on weighting the German survey data in Figure 6 by 0.5. The survey index in East Greenland for 2018 is the average of the 2017 and 2019 values because it was not conducted in 2018.

RESULTS OF THE ASSESSMENT MODEL

Summary of the assessment is shown in Figure 19 and Table 5. The spawning stock increased 1995–2015 but has since then decreased and was on the beginning of 2022 estimated to be close to B_{trigger} . Fishing mortality has been low since 2010, but since the HCR was adopted in 2014, the fishing mortality has been above the target of 0.097 because the catches have exceeded the advice. Recruitment after 2013 is record low for the time series.

Assumptions about the year classes after the 2015 one will not have much effect on the advice this year. This is because the average proportion of fish 10-years old and younger in the landings are only about 10%. Later advice will be affected as well as the development of the spawning stock in short and medium term and is expected to decrease.

Although this year's assessment is consistent with previous assessments it shows a downward revision of SSB and an upward revision of fishing mortality compared to last year's assessment (Figure 20).




Figure 19. Golden redfish. Summary from the assessment 2021. The figure shows total catches, recruitment (age 5) spawning stock biomass (SSB) and fishing mortality for ages 9-19. The dashed line in the SSB plot represents B_{pa} and B_{lim} . The dashed line in the fishing mortality plot indicates the target fishing mortality.




Figure 20. Golden redfish. Comparison of the current assessment (blue line) and the same assessment done in 2018 (red line) and 2019 (green line) for the spawning stock biomass (top), fishing mortality (middle) and recruitment (bottom).

MOHN'S RHO

The evaluation retrospective pattern (five-year peel) of the assessment (Figure 21) is done by calculating the Mohn's rho values. The table below shows the Mohn's rho values for SSB, F and recruitment for five- and ten-year peels:

Variable	Value	
	Five-year peel	Ten-year peel
F_{bar}	-0.0141	-0.0442
SSB	0.00589	0.0231
Rec.	0.704	0.268

The Mohn's rho values for F_{bar} and SSB are low (-1.4% and 0.6% respectively) but indicates that fishing mortality has consistently been underestimated and SSB been overestimated (Figure 21). Mohn's rho for recruitment is on the other hand high (70%) and indicates that recruitment has in previous assessments been overestimated. This value needs though to be taken with caution as recruitment estimates of the five-year peels is very low compared to previous years and any deviation from previous year may have relatively high impact. When extending the peel to 10 years the Mohn's rho value drops to 27%.




Figure 21. Golden redfish. Analytical retrospective pattern of the base run. Recruitment is at age 5 and F shows the development of ages 9–19.

DIAGNOSTICS

Observed and predicted proportion by fleet: Trends in different likelihood components (Figure 22) shows how the fit to survey length distributions has become worse in recent years. This can also be seen in Figure 23 where overall fit to the predicted proportional length distributions in the survey is smaller to the observed for medium sized fish (30–40 cm fish).

Length distributions from the Icelandic commercial catch does usually show good fit except in the most recent period when the large fish is missing and the length distribution narrower (Figure 24).

The fit between predicted and observed age distributions is better than for the length distributions (Figures 25 and 26). The model uses the data as age-length keys in 2 cm intervals for tuning.

Model fit: In Figure 27 the length disaggregated indices are plotted against the predicted numbers in the stock as a time-series. The lack of fit between observed and predicted numbers between 33 and 40 cm is caused by data conflicts with survey indices of larger sizes and compositional data. There appears to be an internal conflict between indices of lengths of 42 cm and above and the large number of smaller fish that was observed in the survey few years earlier. The model results are therefore a compromise between different data sets, and it is not able to follow the amount of 30–40 cm redfish in recent years. The inability of the model to fit the survey biomass in recent years has some support in the characteristics of the survey. Since 2003 most of the biomass in the Icelandic survey has been observed to be aggregated in very dense schools west of Iceland, caught on 5–10 stations every year. The size distribution in those schools is narrow and fish larger than 40 cm were rare.

As the model converges slowly, predicted indices could change several years back when more data are added. However, it is not the magnitude of the residuals but rather the temporal pattern that is worrying (Figure 28). For 35–42 cm fish, the observed indices have been above predictions for 5–11 years. The indices for 41–50 cm fish do not show such temporal pattern although in recent years the observed indices have been below prediction. The correlation between observed and predicted is good for 19–34 cm fish. When looking at the temporal patterns, longevity of the fish must be considered. Positive residuals in size groups 33–38 cm in recent years but negative for most other size groups, especially for fish smaller than 30 cm, indicates narrower length distributions in the survey than predicted (Figure 27).




Figure 22. Golden redfish. Development of component of the objective function with time.




Figure 23. Golden redfish. Fitted proportions-at-length from the Gadget model (black lines) compared to observed proportions in the spring survey (grey lines).




Figure 24. Golden redfish. Fitted proportions-at-length from the Gadget model (black lines) compared to observed proportions from the Icelandic commercial catches (grey lines).




Figure 25. Golden redfish. Fitted proportions-at-age from the Gadget model (black lines) compared to observed proportions in bottom trawl surveys survey (grey lines).




Figure 26. Golden redfish. Fitted proportions-at-age from the Gadget model (black lines) compared to observed proportions from the Icelandic commercial catches (grey lines).




Figure 27. Golden redfish. Gadget fit to disaggregated abundance indices by length from the spring survey.




Figure 28. Golden redfish. Residuals from the fit between model and spring survey indices. The red circles indicate positive residuals (survey results exceed model prediction).

ADVICE

The management plan is based on $F_{9-19} = 0.097$ reducing linearly if the spawning stock is estimated below 220 000 t (B_{trigger}). B_{lim} was proposed as 160 000 t, lowest SSB in the 2012 run. The 2021 SSB was estimated at 260 090 t, and according to the management plan the TAC advice for 2022 was 31 855 t.

REFERENCE POINTS

Harvest control rule (HCR) was evaluated at WKREDMP in January 2014 (ICES, 2014) based on stochastic simulations using the Gadget model. Considering conflicting information by different data continuing for many consequent years, the simulations were conducted using large assessment error with very high autocorrelation ($CV = 0.25$, $\rho = 0.9$).

Yield-per-recruit analysis show that when average size at age 5 was allowed to change after year class 1996, $F_{9-19, \text{MAX}}$ changed from 0.097 to 0.114. The proposed fishing mortality of 0.097 is therefore around 85% of F_{MAX} with current settings. Stochastic simulations indicate that it leads to very low probability of spawning stock going below B_{trigger} and B_{lim} , even with relatively large auto-correlated assessment error.

At WKREDMP 2014, $B_{\text{lim}} = B_{\text{loss}} = 160\,000$ t was defined as the lowest SSB in the 2012 Gadget run. $B_{\text{trigger}} = B_{\text{pa}}$ was defined as 220 000 t by adding a precautionary buffer to the proposed B_{lim} of 160 000 t: $160 \times \exp(0.2 \times 1.645)$. Recruitment in the stochastic simulations was the average of year classes 1975–2003 but those year classes were the basis for the simulations at WKREDMP 2014.

The plot of the average spawning stock against fishing mortality shows that $F_{\text{lim}} = 0.226$ and F_{pa} is then $0.226 / \exp(1.645 \times 0.2) = 0.163$ (Figure 29). The spawning stock decreased considerably from early 1980s to mid-1990s or from 400 000 t to 200 000 t. The reduction in SSB was due to heavy fisheries but SSB increased again gradually because of improved recruitment and lower F .

The probability of current $SSB < B_{trigger}$ is estimated 2.7%. For simplicity, the action of $B_{trigger}$ is not included in the simulations since Gadget is not keeping track of “perceived spawning stock”. Analysis of the stochastic prediction in R shows that if SSB is below $B_{trigger}$ it will only be noted in $<15\%$ of the cases. The reason is that the spawning stock is only likely to go below $B_{trigger}$ in periods of severe overestimation of the stock that occur due to the assumed high autocorrelation in assessment error. This situation differs from that of the stock going below $B_{trigger}$ due to poor recruitment (worse than observed in recent decades). In this case the spawning stock should still have a resilient age structure (as discussed above) and this could reduce the need to take further action below $B_{trigger}$.

Figure 30 shows the development of F_{9-19} based on $F_{9-19} = 0.097$. F is expected to be within the range of the 5th and 95th quantile and the 16th and 84th quantile.




Figure 29. Golden redfish. Average SSB against average fishing mortality and defined reference points.




Figure 30. Golden redfish. Development of F_{9-19} based on $F_{9-19} = 0.097$. The light grey area shows 5th and 95th quantiles and the dark areas 16th and 84th quantiles.

STATE OF THE STOCK

The results from Gadget indicate that fishing mortality has been low since 2009 but above F_{MSY} (Figure 16). Total biomass and SSB have been decreasing since 2016 (Table 5) and the absence of any indications of incoming cohorts raises concerns about the future productivity of the stock.

Results from surveys in Iceland and East Greenland indicate that most recent year classes are poor. The accuracy of the surveys as an indicator of recruitment is not known but recruitment is expected to be poor.

SHORT TERM FORECAST

The Gadget model is length based where growth is modelled based on estimated parameters. The only parameters needed for short term forecast are assumptions about size of those cohorts that have not been seen in the surveys. These year classes were assumed to be the average of the five smallest year classes in 1980–2007.

The results from the short-term simulations based on F_{9-19} is shown in and from short term prognosis with varying fishing mortality in 2022 in Table 6. The results indicate that when fishing according to the management plan the SSB is expected to decrease further and to be below $MSY B_{trigger}$ in 2023 (Table 6).

MEDIUM TERM FORECAST

No medium-term forecast was carried out.

UNCERTAINTIES IN ASSESSMENT AND FORECAST

Various factors regarding the uncertainty and modelling challenges are listed in the WKRED-2012 (ICES, 2012) and WKREDMP-2014 (ICES, 2014) reports.

BASIS FOR ADVICE

Harvest control rule accepted at WKREDMP 2014 (ICES, 2014) and implemented by Icelandic and Greenland authorities in 2014.

MANAGEMENT CONSIDERATION

In 2009 a fishery targeting redfish was initiated in Subarea 14 with annual catches of between 6000 and 8500 t in 2010–2019, highest in 2015 and lowest in 2018. The fishery does not distinguish between species, but based on survey information, golden redfish is estimated to be between 1000 and 2700 in 2010–2015 but increased to 3000–5400 t in 2016–2021.

Subarea 14 is an important nursery area for the entire resource. Measures to protect redfish juveniles in Subarea 14 should be continued (sorting grids in the shrimp fishery).

No formal agreement on the management of *S. norvegicus* exists among the three coastal states, Greenland, Iceland, and the Faroe Islands. However, an agreement was made between Iceland and Greenland in October 2015 on the management of the golden redfish fishery based on the management plan applied in 2014. The agreement was from 2016 to the end of 2018. The agreement states that each year 90% of the TAC is allocated to Iceland and 10% is allocated to Greenland. Furthermore, 350 t are allocated each year to other areas. The plan has not been renewed so no management plan is effective although Iceland and Greenland still follow this plan.

In Greenland and Iceland, the fishery is regulated by a TAC and in the Faroe Islands by effort limitation. The regulation schemes of those states have previously resulted in catches more than TACs advised by ICES.

Since 2009, surveys of redfish in the stock area have consistently shown very low abundance of young redfish (<30 cm). Biomass (SSB and the harvestable biomass) increased from 1995 to 2015 because of recruitment of several strong year classes to the stock. Since then, the biomass has declined. The absence of any indications of any incoming cohorts raises concerns about future productivity of the stock.

ECOSYSTEM CONSIDERATION

Not evaluated for this stock.

REGULATIONS AND THEIR EFFECTS

The separation of golden redfish and Icelandic slope *S. mentella* quota was implemented in the 2010/2011 fishing year.

In the late 1980s, Iceland introduced a sorting grid with a bar spacing of 22 mm in the shrimp fishery to reduce the bycatch of fish juveniles in the shrimp fishery north of Iceland. This was partly done to avoid redfish juveniles as a bycatch in the fishery, but also juveniles of other species. Since the large year classes of golden redfish disappeared out of the shrimp fishing area in the early 1990s, observers report small redfish as being negligible in the Icelandic shrimp fishery. Whether the sorting grids work where the abundance of redfish is high is not known, but not a relevant problem now in 5.b as abundance of small redfish is low and shrimp fisheries limited.

There is no minimum landing size of golden redfish in Division 5.a. However, if more than 20% of a catch observed on board is below 33 cm a small area can be closed temporarily. A large area west and southwest of Iceland is closed permanently for fishing to protect young golden redfish.

There is no regulation of the golden redfish in Division 5.b.

Since 2002 it has been mandatory in the shrimp fishery in Subarea 14 to use sorting grids to reduce bycatches of juvenile redfish in the shrimp fishery.

CHANGES IN FISHING TECHNOLOGY AND FISHING PATTERNS

There have been no changes in the fishing technology and the fishing pattern of golden redfish in ICES Subareas 5 and 14.

BENCHMARK

Benchmark meeting for golden redfish is scheduled in 2023.

Golden redfish was last benchmarked in 2014 and the group thinks that benchmarking the stock is of high importance. The proposed benchmark meeting will explore several issues of current assessment model. These include poor fit to survey indices for fish between 30–40 cm; potential dome-shape in selectivity; uncertainty estimates are not available; investigate the appropriateness of the current growth and maturity model used in the assessment. In addition, the meeting will explore alternative assessment methods. Underutilized data sources from ICES 5.b and 14.b, mainly relevant survey and commercial samples of age and length. Biological reference points will need to be redefined depending on the assessment method, especially in relation to the $F_{0.5}$. Change in form of harvest control rule will also be explored, that is change the rule to proportion of biomass above certain size (i.e., 33 cm and bigger fish) from the F -based rule that is used now.

REFERENCES

- ICES 2012. Report of the Benchmark Workshop on Redfish (WKRED 2012). ICES CM 2012/ACOM:48, 291 pp.
- ICES 2014. Report of the Workshop on Redfish Management Plan Evaluation (WKREDMP). ICES CM 2014/ACOM:52, 269 pp.
- Pálsson, Ó., Björnsson, H., Björnsson, E., Jóhannesson, G. and Ottesen P. 2010. Discards in demersal Icelandic fisheries 2009. Marine Research in Iceland 154.

TABLES

Table 1. Survey indices and CV of golden redfish from the spring survey 1985–2021 and the autumn survey 1996–2020.

YEAR	SPRING SURVEY		AUTUMN SURVEY	
	BIOMASS	CV	BIOMASS	CV
1985	307,926	0.095		
1986	327,765	0.120		
1987	322,121	0.122		
1988	253,559	0.095		
1989	281,117	0.122		
1990	242,450	0.223		
1991	199,128	0.114		
1992	160,545	0.088		
1993	179,275	0.130		
1994	171,135	0.097		
1995	146,102	0.102		
1996	195,697	0.164	199,793	0.248
1997	212,558	0.216	120,628	0.279
1998	206,461	0.136	186,505	0.348
1999	297,090	0.143	262,691	0.310
2000	221,279	0.176	141,940	0.200
2001	192,724	0.176	177,456	0.155
2002	250,420	0.173	192,813	0.150
2003	333,901	0.161	199,450	0.159
2004	326,868	0.236	220,308	0.241
2005	310,635	0.129	229,013	0.240
2006	257,010	0.157	279,290	0.335
2007	339,778	0.224	219,951	0.252
2008	247,895	0.154	288,149	0.244
2009	302,204	0.253	294,028	0.282
2010	383,407	0.245	227,335	0.171
2011	401,358	0.235		
2012	461,921	0.204	343,115	0.225
2013	457,451	0.177	317,325	0.156
2014	402,773	0.174	431,369	0.232
2015	406,150	0.281	360,722	0.173
2016	615,712	0.313	401,135	0.279
2017	507,058	0.205	428,351	0.187
2018	497,092	0.210	342,467	0.195
2019	410,550	0.158	383,532	0.233
2020	411,320	0.206	244,099	0.159
2021	441,154	0.194	269,053	0.199
2022	378,907	0.177		

Table 2. Golden redfish in 5.a. Age disaggregated indices (in millions) from the autumn groundfish survey 1996–2020. The survey was not conducted in 2011.

YEAR/AGE	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
1	0.3	1.0	3.6	3.3	0.8	0.4	0.1	0.0	0.0	0.1	0.2	0.1	0.0	0.1	0.0		0.0	0.0	0.0
2	2.4	0.2	1.5	3.3	1.7	1.0	0.9	0.5	0.2	0.1	0.6	1.2	0.3	0.3	0.0		0.0	0.0	0.2
3	0.7	2.2	0.9	3.3	1.4	1.9	1.5	1.1	1.0	0.2	0.7	1.2	2.5	0.4	1.7		0.1	0.0	0.3
4	1.6	1.6	2.3	1.5	1.6	2.4	6.1	1.1	1.8	1.0	0.5	1.1	2.7	4.4	0.3		1.4	0.2	0.1
5	8.3	2.2	0.9	4.7	1.2	5.4	5.8	12.3	3.3	4.2	5.0	2.1	4.1	12.0	4.3		4.1	1.0	0.8
6	40.0	6.9	3.5	2.8	7.9	2.1	11.8	17.7	28.6	4.8	6.8	10.4	7.9	11.6	14.2		3.1	4.1	1.8
7	11.3	22.5	16.6	10.5	6.7	10.8	3.3	38.2	36.7	39.7	15.6	26.0	39.2	13.9	15.1		23.5	3.0	12.8
8	19.1	14.3	58.2	47.2	6.4	10.9	26.9	9.9	65.4	44.9	81.9	35.8	75.1	73.9	23.4		70.3	41.8	24.6
9	15.1	13.0	22.4	99.9	26.2	7.1	11.2	48.5	21.0	62.7	81.5	76.6	67.9	96.4	54.4		60.6	84.8	96.9
10	28.9	11.1	26.1	43.7	95.0	17.3	16.6	12.7	45.6	24.9	85.7	37.4	106.4	58.7	69.0		62.9	56.3	151.8
11	102.7	17.6	18.9	20.7	11.5	111.2	32.0	17.0	19.3	44.2	26.3	36.1	63.2	100.9	32.5		103.8	41.3	90.8
12	16.2	67.8	19.1	16.8	14.2	23.6	116.3	39.7	13.4	19.6	37.5	19.0	55.1	45.9	57.4		74.2	68.6	69.7
13	10.1	6.2	104.5	20.8	7.9	23.6	20.0	111.3	26.6	15.4	18.0	23.8	13.5	42.9	28.6		43.3	47.5	67.5
14	16.8	5.3	10.1	147.1	8.0	7.9	11.5	12.4	103.9	26.8	15.1	8.2	18.2	10.2	19.6		39.1	26.5	50.4
15	33.9	7.2	7.6	6.0	51.4	9.2	9.8	10.8	13.6	82.1	18.3	6.8	9.1	18.3	9.1		19.6	31.7	27.0
16	16.1	10.0	7.8	9.6	5.3	58.9	10.4	6.1	9.6	9.5	75.4	16.9	7.8	6.9	10.9		16.7	18.7	26.6
17	1.9	6.9	14.1	10.9	2.5	4.3	45.4	7.5	6.0	6.7	8.7	49.4	13.1	6.4	4.7		6.1	12.8	17.1
18	1.7	3.9	7.6	11.1	2.5	5.0	4.6	32.7	6.1	3.7	4.3	10.4	36.6	7.4	3.1		5.9	7.2	12.3
19	4.3	2.0	0.5	8.4	4.6	3.6	3.0	4.5	21.6	5.0	2.8	4.5	6.2	28.4	6.6		3.9	5.2	6.0
20	6.6	1.4	3.2	3.9	6.5	4.1	3.2	1.6	3.1	22.0	3.1	1.5	5.7	4.7	22.2		3.9	4.5	5.9
21	1.1	0.8	2.3	2.8	1.0	3.7	3.9	1.1	1.8	2.5	17.8	4.0	2.1	2.1	3.1		3.5	4.8	4.8
22	5.0	1.5	0.8	1.0	1.6	2.3	3.2	2.7	1.7	2.1	2.0	13.8	2.3	1.3	1.2		18.3	2.4	3.6
23	3.9	2.4	2.2	2.1	0.4	0.3	0.8	1.1	2.5	2.4	1.7	1.3	11.0	2.0	1.6		2.9	18.2	3.4
24	4.6	0.8	0.4	0.6	1.0	0.5	0.4	0.3	0.0	0.9	1.0	1.3	1.4	10.2	0.7		2.0	2.6	12.7
25	3.9	2.7	1.4	2.8	0.8	0.3	0.5	0.3	1.2	1.2	1.7	0.2	0.8	0.8	5.7		1.2	1.2	1.5
26	0.9	1.1	0.2	1.2	0.7	0.5	0.6	0.2	0.4	0.3	0.9	0.6	0.9	1.0	0.6		1.7	1.1	0.9
27	0.9	0.2	0.9	2.9	0.5	0.8	0.3	0.3	0.0	0.1	0.9	0.3	1.2	1.3	0.4		7.5	0.8	0.9
28	0.8	0.4	0.5	1.5	0.7	0.5	0.2	0.0	0.2	0.2	0.2	0.0	0.6	0.2	0.7		0.4	8.7	0.5
29	0.1	0.0	0.5	1.2	0.5	0.2	0.7	0.1	0.2	0.0	0.4	0.4	0.8	1.6	0.4		0.4	0.5	3.3
30+	0.8	1.4	3.0	1.1	1.3	2.3	1.7	1.5	1.6	2.1	1.0	0.9	1.5	1.7	2.0		2.1	3.5	2.6
TOTAL	360.0	214.6	341.6	492.7	271.8	322.1	352.7	393.2	436.4	429.4	515.6	391.3	557.2	565.9	393.5		582.5	499.2	696.9

Table 4. Golden redfish in 5.a. Continued.

2015	2016	2017	2018	2019	2020	2021
0.0	0.0	0.0	0.1	0	0.4	0.3
0.1	0.0	0.3	0.2	0.1	0.2	0.2
0.6	0.0	0.3	0.4	0.4	1.0	0.2
0.3	1.8	0.2	0.1	0.8	0.7	0.6
0.1	0.3	1.6	0.2	1.5	1.3	1.3
1.2	0.8	1.3	3.0	0.9	0.8	2.5
7.6	3.9	1.6	2.5	15.3	0.7	1.3
28.3	29.1	10.4	2.0	7.8	10.9	1.6
33.1	63.8	38.1	5.9	7.4	3.9	12.4
86.4	48.1	93.8	36.7	20.3	7.4	7.0
100.7	87.5	56.9	72.1	46.8	18.4	9.0
52.9	97.2	95.7	58.4	91.5	41.0	30.4
47.6	54.3	87.8	65.7	58.7	39.1	35.9
41.7	45.3	41.9	54.9	62.7	24.3	48.7
40.3	35.8	27.4	27.3	45.4	39.0	14.9
21.1	31.9	28.8	20.2	36.1	25.7	36.4
20.0	20.3	35.6	21.9	18.7	10.5	23.2
10.0	22.1	17.8	21.1	21.7	12.1	13.1
10.0	16.1	14.7	12.9	22.1	12.0	10.3
9.9	8.9	16.8	11.3	13.7	11.1	10.8
3.3	3.0	11.5	6.0	14.7	6.9	12.4
2.5	3.9	4.8	10.3	12.3	4.6	9.2
2.1	3.7	6.1	6.9	7.2	4.1	8.4
1.1	2.8	4.8	2.8	3.7	3.3	5.6
13.1	3.4	2.9	2.6	1.3	2.5	4.4
1.5	15.0	2.6	2.9	2.0	1.8	2.7
1.4	1.0	13.9	2.6	1.3	1.9	1.5
1.6	1.0	1.7	11.5	1.7	0.8	0.8
1.0	0.9	1.8	1.5	10.4	1.3	2.7
6.9	6.7	7.9	7.5	5.3	9.6	14.8
546.3	608.9	629.0	472.0	531.8	297.4	322.6

Table 3. Official landings (in tonnes) of golden redfish, by area, 1978–2020 as officially reported to ICES.

YEAR	AREA		ICES 6	ICES 14	TOTAL
	ICES 5.A	ICES 5.B			
1978	31 300	2 039	313	15 477	49 129
1979	56 616	4 805	6	15 787	77 214
1980	62 052	4 920	2	22 203	89 177
1981	75 828	2 538	3	23 608	101 977
1982	97 899	1 810	28	30 692	130 429
1983	87 412	3 394	60	15 636	106 502
1984	84 766	6 228	86	5 040	96 120
1985	67 312	9 194	245	2 117	78 868
1986	67 772	6 300	288	2 988	77 348
1987	69 212	6 143	576	1 196	77 127
1988	80 472	5 020	533	3 964	89 989
1989	51 852	4 140	373	685	57 050
1990	63 156	2 407	382	687	66 632
1991	49 677	2 140	292	4 255	56 364
1992	51 464	3 460	40	746	55 710
1993	45 890	2 621	101	1 738	50 350
1994	38 669	2 274	129	1 443	42 515
1995	41 516	2 581	606	62	44 765
1996	33 558	2 316	664	59	36 597
1997	36 342	2 839	542	37	39 761
1998	36 771	2 565	379	109	39 825
1999	39 824	1 436	773	7	42 040
2000	41 187	1 498	776	89	43 550
2001	35 067	1 631	535	93	37 326
2002	48 570	1 941	392	189	51 092
2003	36 577	1 459	968	215	39 220
2004	31 686	1 139	519	107	33 451
2005	42 593	2 484	137	115	45 329
2006	41 521	656	0	34	42 211
2007	38 364	689	0	83	39 134
2008	45 538	569	64	80	46 251
2009	38 442	462	50	224	39 177
2010	36 155	620	220	1 653	38 648
2011	43 773	493	83	1 005	45 354
2012	43 089	491	41	2 017	45 635
2013	51 330	372	92	1 499	53 263
2014	47 769	201	60	2 706	50 736
2015	48 769	270	44	2 562	51 645
2016	54 041	165	50	5 442	59 698
2017	50 119	1 418	93	4 501	56 141
2018	48 014	1 129	80	4 004	53 227
2019	44 746	1 119	101	2 665	48 530
2020	40 688	1 304	100	4 105	46 197
2021 ¹⁾	39 616	178	100	3 532	43 426

1) Provisional

Table 4. Golden redfish in 5.a. Observed catch in weight (tonnes) by age and years in 1995–2021.

YEAR/AGE	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
7	46	0	33	24	6	38	125	127	191	226	227	176	135	215	103	60	138	68	30	235
8	321	389	226	280	342	62	143	884	201	855	755	987	446	1,057	936	359	558	612	555	475
9	1,432	867	481	586	1,592	825	402	736	1,312	501	1,877	2,134	1,727	2,164	1,689	2,218	1,626	1,603	2,197	1,752
10	8,598	3,887	1,039	1,193	1,252	4,180	1,653	808	1,080	2,107	1,496	3,605	2,442	5,006	3,059	2,725	4,772	3,444	3,886	6,176
11	2,570	9,575	2,708	1,118	1,843	1,843	7,768	3,192	1,160	828	3,093	2,017	3,319	3,997	4,964	2,786	5,699	6,725	5,952	6,751
12	1,286	2,170	11,609	3,221	2,521	2,224	1,810	10,955	3,863	989	1,899	2,789	1,911	4,682	4,457	4,921	4,899	7,345	9,488	5,807
13	3,616	1,354	2,828	12,425	2,447	1,665	1,930	3,012	9,576	2,017	1,366	1,624	3,068	2,297	3,430	3,895	6,235	4,021	6,896	5,809
14	5,787	1,523	1,366	2,068	15,536	2,329	1,243	2,548	2,304	8,612	3,021	1,275	1,050	2,819	1,848	2,740	3,772	4,721	4,032	4,776
15	6,229	4,293	3,106	2,020	1,242	14,598	826	1,805	1,932	2,148	11,840	2,818	955	1,546	2,008	1,378	2,501	2,668	4,466	3,061
16	1,833	5,033	3,579	2,394	1,250	1,752	11,487	2,998	1,202	1,656	2,073	10,318	2,168	1,067	1,247	1,201	1,309	1,525	3,043	2,538
17	912	954	2,968	3,404	1,795	1,170	515	11,726	2,231	870	1,447	2,074	9,337	1,804	681	820	981	820	1,720	1,921
18	395	372	869	2,029	2,619	1,602	769	2,054	6,494	1,381	1,243	1,191	1,329	8,188	1,502	648	602	813	1,205	1,245
19	1,244	252	616	1,013	2,194	2,400	1,025	1,150	784	5,065	1,241	722	741	1,503	6,158	1,086	691	492	764	464
20	1,232	343	919	723	1,237	2,141	1,684	622	390	1,093	6,387	956	717	966	970	4,980	987	808	488	1,202
21	549	1,059	440	528	452	538	916	1,360	585	342	387	5,524	876	567	654	901	5,052	627	510	438
22	674	698	534	397	211	438	386	982	840	464	456	552	4,765	831	576	762	1,056	3,512	772	425
23	1,521	790	641	426	326	283	399	697	788	599	758	226	732	4,231	342	519	753	477	3,298	486
24	695	0	567	660	215	63	155	352	426	528	591	396	113	382	2,561	665	204	324	183	2,929
25	777	0	703	536	810	408	119	270	307	239	417	457	599	254	98	2,151	134	225	199	183
26	396	0	263	382	264	361	109	176	71	94	94	97	329	433	97	199	1,336	237	171	195
27	372	0	135	432	592	220	265	80	74	187	253	254	345	337	199	348	77	1,326	108	142
28	799	0	186	358	227	520	182	287	26	123	161	200	199	169	94	131	201	198	918	57
29	0	0	137	54	105	379	142	469	95	127	28	168	36	171	359	155	44	72	37	674
30+	230	0	388	501	745	1,152	1,015	1,280	643	636	1,484	962	1,024	851	411	507	145	426	414	33
TOTAL	41,515	33,558	36,339	36,771	39,823	41,188	35,066	48,569	36,576	31,688	42,591	41,520	38,364	45,537	38,443	36,156	43,773	43,088	51,328	47,768

Table 4. Golden redfish in 5.a. Continued.

YEAR/AGE	2015	2016	2017	2018	2019	2020	2021
7	14	49	0	0	214	0	41
8	563	751	104	51	144	507	26
9	902	2,717	949	212	64	288	1,276
10	3,154	3,713	4,503	2,279	1,227	575	766
11	7,118	8,111	3,523	4,890	4,678	2,185	373
12	7,104	9,393	7,077	4,812	6,176	4,928	2,440
13	5,553	6,688	8,748	6,507	4,028	4,154	4,056
14	5,673	4,705	5,370	7,779	5,710	3,148	4,743
15	4,774	4,024	3,790	4,278	5,127	8,115	3,794
16	3,015	2,629	3,576	3,243	4,006	5,032	5,350
17	2,651	2,729	3,012	2,748	2,607	2,253	4,801
18	1,861	2,013	1,866	2,614	2,301	1,545	2,310
19	780	1,724	1,412	1,282	1,376	1,329	1,167
20	1,192	663	1,187	1,347	1,512	1,564	1,646
21	288	536	990	1,211	1,147	788	1,261
22	275	350	438	629	508	970	768
23	196	223	489	496	518	522	942
24	424	241	313	277	161	600	799
25	1,816	304	324	336	56	82	152
26	243	1,335	148	167	184	45	443
27	214	176	1,265	35	350	62	28
28	189	29	87	1,663	103	122	186
29	87	25	192	26	1,161	162	214
30+	682	907	756	1,133	1,387	1,713	2,030
TOTAL	48,770	54,043	50,117	48,015	44,745	40,689	39,616

Table 5. Golden redfish. Results from the Gadget model of total biomass, spawning stock biomass, recruitment at age 5 (in millions), catch and fishing mortality, projections are in *italic*. All weights are in tonnes.

YEAR	BIOMASS	SSB	R(AGE5)	CATCHES	F9-19
1971	534,085	338,242	210.6	67,880	0.115
1972	532,312	326,889	161.7	50,890	0.089
1973	545,411	331,438	456.7	43,719	0.076
1974	604,345	347,290	220.5	50,598	0.083
1975	639,480	362,455	117.9	61,920	0.097
1976	654,476	373,340	195.9	94,420	0.145
1977	645,558	361,350	191.0	53,753	0.087
1978	678,240	389,424	129.6	48,736	0.071
1979	709,833	425,542	165.5	77,212	0.106
1980	715,593	439,297	100.2	89,143	0.120
1981	699,854	441,430	86.9	101,966	0.142
1982	665,192	429,215	64.2	130,322	0.193
1983	593,378	386,355	66.1	106,050	0.171
1984	540,066	357,097	71.8	95,288	0.163
1985	493,226	330,502	129.5	78,531	0.138
1986	466,558	312,661	123.3	76,908	0.146
1987	439,463	291,881	63.4	76,559	0.158
1988	404,205	266,908	39.1	89,804	0.212
1989	349,457	226,159	42.5	56,645	0.150
1990	325,284	211,258	347.6	66,314	0.199
1991	326,320	187,851	57.5	56,015	0.186
1992	309,854	172,180	38.6	55,826	0.206
1993	289,910	156,762	52.1	50,179	0.205
1994	275,205	147,027	61.7	42,520	0.183
1995	267,797	144,733	325.4	44,263	0.195
1996	289,841	143,002	83.5	35,595	0.155
1997	298,610	149,041	38.5	38,996	0.165
1998	298,230	152,952	38.8	39,694	0.165
1999	295,290	157,102	76.5	42,463	0.175
2000	291,814	159,403	47.5	42,607	0.172
2001	283,408	160,947	103.2	36,744	0.142
2002	290,677	167,273	111.8	50,730	0.195
2003	285,287	161,447	163.6	38,219	0.148
2004	301,906	166,336	101.6	32,766	0.123
2005	316,429	176,691	154.3	46,619	0.173
2006	326,245	177,413	151.7	42,108	0.161
2007	347,359	185,875	98.4	39,154	0.146
2008	362,743	199,321	119.8	46,195	0.165
2009	375,258	210,288	184.1	39,301	0.133
2010	408,289	230,741	155.3	38,504	0.119
2011	439,377	254,843	83.2	45,146	0.130
2012	450,600	274,185	123.6	45,423	0.123
2013	468,061	293,834	76.1	53,223	0.137
2014	467,648	305,377	36.3	50,697	0.126
2015	459,245	315,360	11.0	51,621	0.124
2016	440,842	319,575	13.3	59,711	0.142
2017	409,897	310,673	35.9	56,355	0.136
2018	382,627	298,542	4.5	53,167	0.133
2019	349,029	282,338	8.2	48,550	0.128
2020	317,118	264,207	19.1	46,116	0.129
2021	286,687	242,926	26.2	43,337	0.134
2022	258,329	220,056	47.2		

Table 6. Golden redfish. Output from short term prognosis. Multiplier is based on reference to the adopted HCR $F_{9-19} = 0.097$. All weights are in tonnes.

Biomass (2022)	SSB (2022)	F9-19 (2022)	Landings (2022)	Biomass (2023)	SSB (2023)
258 329	220 056	0.128	37 241	238 910	200 045

Basis	Total catch (2023)	F9-19 (2023)	Biomass 5+ (2024)	SSB (2024)
Management plan	25 545	0.097	229 871	189 588
Other catch options				
F_0	0	0	255 771	213 812
$F_{sq} = F_{2021}$	31 152	0.134	224 183	184 271