

WHITING - LÝSA

Merlangius merlangus

GENERAL INFORMATION

Whiting is a demersal gadoid species like cod and haddock, but smaller with a maximum length of about 80 cm, males and females being similar in size. In Icelandic waters, sexual maturity is reached at around 30 cm.

THE FISHERY

Whiting has been caught mainly as bycatch all around Iceland in recent years, but mostly around south and west of Iceland (Figures 1 and 2). Annual catches have been between 500 and 1000 tonnes except for 2008-2012 when catches peaked in 2011 and were 2602 tonnes (Figure 2). Increased catches in this period occurred almost exclusively in the southwest (Figure 2). Whiting is found at depths ranging from 10 to 300 m but is mostly caught between 100 and 250 m (Figure 3).

Whiting is mainly caught in demersal trawls but to some extent in *Nephrops* trawls, longline and demersal seine (Table 1, Figure 4). The number of boats reporting whiting catches increased with increased catches between 2007 and 2012 but has since then decreased (Figures 2 and 4, and Table 1).




Figure 1. Whiting. Geographic distribution of the Icelandic fishery 2012-2020 as reported in logbooks.




Figure 2. Whiting. Catch distribution and proportions by area 2000-2020 according to logbooks.




Figure 3. Whiting. Depth distribution of catches 2000-2020 according to logbooks.




Figure 4. Whiting. Total catch (landings) of whiting by fishing gear 1994-2020 according to statistics from the Directorate of Fisheries.

Table 1. Whiting. Number of Icelandic boats reporting catches of whiting, landings by fishing gear and yearly reported landings according to the Directorate of Fisheries.

YEAR	NUMBER OF VESSELS				CATCHES (TONNES)				
	<i>Demersal trawl</i>	<i>Nephrops trawl</i>	<i>Longline</i>	<i>Other</i>	<i>Demersal trawl</i>	<i>Nephrops trawl</i>	<i>Longline</i>	<i>Other</i>	<i>Sum</i>
2000	76	13	131	79	1037	51	157	99	1344
2001	59	15	111	102	792	136	114	131	1173
2002	62	23	81	92	913	150	90	145	1298
2003	54	22	100	94	671	62	153	139	1025
2004	51	18	116	76	682	22	224	110	1038
2005	54	13	115	77	488	26	205	79	798
2006	50	15	144	83	439	29	460	124	1052
2007	53	7	181	90	741	22	394	102	1259
2008	58	12	190	84	928	21	557	182	1688
2009	56	13	201	151	1404	35	520	349	2308
2010	52	17	186	133	2036	155	425	234	2850
2011	52	15	187	120	2288	176	345	156	2965
2012	46	15	174	102	777	178	320	194	1469
2013	37	15	172	70	417	240	255	70	982
2014	33	15	154	69	518	124	205	81	928
2015	32	13	130	41	546	129	115	69	859
2016	36	11	127	33	494	65	103	71	733
2017	27	8	95	23	360	41	107	44	552
2018	32	8	72	28	659	30	60	68	817
2019	39	8	76	28	581	25	93	62	761
2020	39	5	72	29	529	19	67	19	634

LENGTH DISTRIBUTIONS FROM COMMERCIAL CATCHES OF WHITING

Length measurements of whiting from commercial catches are scarce and missing for some years. Most whiting caught in the commercial fishery are 38-55 cm (Figure 6).




Figure 6. Whiting. Relative length distribution from commercial catches in 1996-2019, excluding years where measurements were lacking.

SURVEY DATA

Annual Icelandic groundfish surveys have been conducted in March (IS-SMB) since 1985 and October (IS-SMH) since 1996. Both surveys cover the distribution area of whiting on Icelandic grounds. For monitoring, harvestable biomass and recruitment indices were estimated for both surveys (Figure 7). The harvestable biomass index is calculated as the biomass of individuals 40 cm and larger. The recruitment index is defined as whiting smaller than 20 cm.

Both the total biomass index and harvestable biomass index in IS-SMB increased from 2003 to a maximum in 2005 but decreased to a low level in 2015 (Figure 7). Since then, both indices have increased and are now close to the mean of the time series (1985-2020). The biomass indices from IS-SMH are much more variable but show similar trends in the last decade. Recruitment indices show similar trends in both surveys (Figure 7). Strong recruitment was observed in 2003, 2007 and 2020 in IS-SMH and in 2004, 2008 and 2021 in IS-SMB. These peaks can be seen in the length distributions (Figures 8 and 9) and reached the harvestable biomass 2-3 years later. In the past five years, a slight increase in recruitment was observed with a substantial increase in the past year. In concordance with recruitment, total- and harvestable biomass have been increasing in both IS-SMB and IS-SMH.

Spatial distribution of whiting from the spring survey is similar to what is observed in the commercial catches, that is, mostly in the southwest of Iceland (Figures 1 and 2, 10 and 11). The autumn survey however shows the highest indices in a larger area, southeast, southwest, and west (Figures 12 and 13).




Figure 7. Whiting. Total biomass indices (upper left), harvestable biomass indices (≥ 40 cm, upper right), and juvenile abundance indices (≤ 20 cm, lower) from IS-SMB (blue) 1985-2021 and IS-SMH (red) 1996-2020, along with 95% CI.




Figure 8. Whiting. Length-disaggregated abundance indices from IS-SMB 1985-2021. The blue line shows the mean for all years. Note different scales on y-axes.




Figure 9. Whiting. Length-disaggregated abundance indices from IS-SMH 1996-2020. The blue line shows the mean for all years. Note different scales on y-axes.




Figure 10. Whiting. Spatial distribution from IS-SMB in 2021.




Figure 11. Whiting. Spatial distribution of biomass index from IS-SMB 1985-2021.




Figure 12. Whiting. Spatial distribution of catches from IS-SMH 2020.




Figure 13. Whiting. Spatial distribution of biomass index from IS-SMH 1996-2020.

MANAGEMENT

Whiting has not been subject to management such as TAC limitations and hence, catch advice has not been given by the Marine and Freshwater Research Institute before 2019.

Table 2. Whiting. Recommended TAC, national TAC set by the Ministry, and landings (tonnes).

Fishing year	Rec. Tac	National TAC	Catch
2001/02	-	-	1192
2002/03	-	-	1309
2003/04	-	-	1001
2004/05	-	-	964
2005/06	-	-	895
2006/07	-	-	1030
2007/08	-	-	1812
2008/09	-	-	1984
2009/10	-	-	2835
2010/11	-	-	3249
2011/12	-	-	1601
2012/13	-	-	1060
2013/14	-	-	1034
2014/15	-	-	877
2015/16	-	-	690
2016/17	-	-	642
2017/18	-	-	844
2018/19	-	-	780
2019/20	836	-	607
2020/21	1003	-	